arXiv:1607.06418v1 [math.GR] 21 Jul 2016

July 22, 2016

GROUP BASES FOR SOME SOLVABLE GROUPS AND SEMIDIRECT PRODUCTS

BRET BENESH AND JASON LUTZ

Abstract. A set B is a basis for a vector space V if every element of V can be uniquely written as a linear combination of the elements of B. There is a similar definition of a basis for a finite group. We show that certain semidirect products of finite groups—including all semidirect products of finite abelian groups—have bases; any group of order m or $2m$ for odd, cube-free m has a basis; and the quaternions do not have a basis.

1. Introduction and Preliminaries

Let G be a finite group, and suppose there exists $(b_1, \dots, b_n) \in (G \setminus \{e\})^n$ such that for every $g \in G$, there are unique $a_i \in \{0, \dots, |b_i| - 1\}$ such that $g = b_1^{a_1} \cdots b_n^{a_n}$. In this case, we say that $(b_1, \dots, b_n)$ is a *basis* for G and $(a_1, \dots, a_n)$ is a *representation* for g, and it is easy to see that $|b_1| \cdots |b_n| = |G|$. If $G = \{e\}$, we say that G has a *trivial basis*.

Philip Hall [3] introduced the idea of a group basis, and Charles Hopkins [4] proved that dihedral, symmetric, and alternating groups all have bases. There are several other families of groups that have natural bases. If $G = \langle b \rangle$ is cyclic, then (b) is easily seen to be a basis of G. The Fundamental Theorem of Abelian Groups states that if A is a finite abelian group, then there are $b_i \in A$ and prime powers q_i such that $A \cong \langle b_1 \rangle \times \dots \langle b_m \rangle$, where $|b_i| = q_i$. It is easy then to see that $(b_1, \dots, b_m)$ is a basis for A. Finally, groups can have more than one basis: $((1, 1))$ and $((1, 0), (0, 1))$ are both bases for $\mathbb{Z}_2 \times \mathbb{Z}_3$.

A group basis is a special case of the following definition by Magliveras [7]: Let G be a finite group, and let α be a sequence $(\alpha_1, \dots, \alpha_n)$, where each α_i is itself a sequence $(a_{i,1}, \dots, a_{i,r_i})$ of elements of G. Then α is said to be a *logarithmic signature* for G if $|G| = r_1 \cdots r_n$ and each $g \in G$ is uniquely represented as a product $g = a_{1,j_1} \cdots a_{n,j_n}$ with $a_{i,j_i} \in \alpha_i$. Logarithmic signatures have been considered for cryptographic systems in [6], [8], and [9]. Then a basis for a finite group G is a logarithmic signature $(\alpha_1, \dots, \alpha_n)$ such that each α_i is a cyclic subgroup of G. All solvable groups have a logarithmic signature [2]. For instance, recall that the quaternions Q_8 is the set $\{\pm 1, \pm i, \pm j, \pm k\}$, where $|1| = 1$, $|-1| = 2$ and $|\pm x| = 4$ for $x \in \{i, j, k\}$. One such logarithmic signature for Q_8 is $((1, i, -1, -i), (1, j))$. However, this logarithmic signature is not a basis because $\{1, j\}$ is not equal to a cyclic subgroup of Q_8. In fact, the proposition below demonstrates that Q_8 has no basis, so we say that that Q_8 is *basisless*.

Proposition 1.1. *The quaternions Q_8 do not have a basis.*

Proof. Suppose Q_8 has a basis $(b_1, \dots, b_n)$. Then $|b_1| \cdots |b_n| = 8$, and we see that $|b_i| \in \{2, 4, 8\}$ with $1 \leq n \leq 3$. Since Q_8 is noncyclic, we can refine this to $|b_i| \in \{2, 4\}$ and $n \in \{2, 3\}$. Since Q_8 has a unique element -1 of order 2, we conclude that the basis is either $(-1, b)$ or $(b, -1)$ for some $b \in Q_8$ with $|b| = 4$. Then b^2 has order 2, so $b^2 = -1$. If the basis is $(-1, b)$, we then have $(0, 2)$ and $(1, 0)$ are both representations for -1, which contradicts the uniqueness of representations of the basis. Similarly, $(0, 1)$ and $(2, 0)$ are representations for -1 for the basis $(b, -1)$, which is also a contradiction. Therefore, no basis can exist for Q_8. □

Groups involving Q_8 can still have a basis. For instance, the semidihedral group of order 16 with presentation $\langle a, x \mid a^8 = x^2 = e, x^{-1}ax = a^{-1} \rangle$ has a normal subgroup $\langle a^2, ax \rangle$ isomorphic to the quaternions, yet has a basis (a, x). Also, there is a group G of order 16 with presentation $\langle a, x \mid a^4 = x^4 = e, x^{-1}ax = a^{-1} \rangle$ such that $G/\langle a^2x^2 \rangle \cong Q_8$, yet G has a basis (a, x) (Corollary 2.2 also demonstrates that G has a basis, since G is isomorphic to the semidirect product $\mathbb{Z}_4 \rtimes \mathbb{Z}_4$ with an inversion action). Finally, the symmetric group S_8 has a non-normal subgroup isomorphic to the quaternions, and S_8 has a basis [4]. So it is possible for a group with a basisless subgroup (normal or otherwise) or basisless quotient to have a basis.

We end this section with an important lemma.

Lemma 1.2. *Let G be a finite group, and let H be a subgroup of G with a basis $(h_1, \dots, h_m)$. Suppose there exists $b_1, \dots, b_n \in G \setminus H$ such that, for all $x = b_1^{c_1} \cdots b_n^{c_n}$ and $y = b_1^{k_1} \cdots b_n^{k_n}$ with $c_i, k_i \in \{0, \dots, |b_i| - 1\}$, the following hold:*

1. *$x = y$ if and only if $(c_1, \dots, c_n) = (k_1, \dots, k_n)$,*
2. *$xy^{-1} \in H$ if and only if $x = y$, and*
3. *$|G : H| = |b_1| \cdots |b_n|$.*

Then $B = (h_1, \dots, h_m, b_1, \dots, b_n)$ is a basis for G.

Proof. We start by showing that every coset of H can be written as $Hb_1^{c_1} \cdots b_n^{c_n}$ for some $(c_1, \dots, c_n) \in \prod_{i=1}^n \{0, \dots, |b_i| - 1\}$. Let ψ be a map from $\prod_{i=1}^n \{0, \dots, |b_i| - 1\}$ to the set of right cosets of H defined by $\psi(c_1, \dots, c_n) = Hb_1^{c_1} \cdots b_n^{c_n}$. Then ψ is injective by Items (1) and (2) and surjective by Item (3), giving the result.

Let $g \in G$. Then g appears in exactly one coset of H, so let $(c_1, \dots, c_n)$ be such that $g \in Hb_1^{c_1} \cdots b_n^{c_n}$. Then $g = hb_1^{c_1} \cdots b_n^{c_n}$ for some $h \in H$, and we can write h uniquely as $h_1^{l_1} \cdots h_m^{l_m}$ for some $(l_1, \dots, l_m)$. Then

$$g = h_1^{l_1} \cdots h_m^{l_m} \cdot b_1^{c_1} \cdots b_n^{c_n}$$

for unique $l_i \in \{0, \dots, |h_i| - 1\}$ and unique $c_i \in \{0, \dots, |b_i| - 1\}$. Therefore, $(h_1, \dots, h_m, b_1, \dots, b_n)$ is a basis for G. □

2. Families of groups

2.1. **Semidirect products of abelian groups.** Let H and N be finite groups such that there is a map $\phi : H \to \mathrm{Aut}(N)$, and let $\phi_h : N \to N$ denote $\phi(h)$. Recall that a *semidirect product* $N \rtimes_\phi H$ is defined to be the set $\{(h, n) \mid h \in H, n \in N\}$ together with the operation defined by $(h_1, n_1)(h_2, n_2) = (h_1 h_2, \phi_{h_2}(n_1) n_2)$ for all $h_i \in H$ and $n_i \in N$.

The following results generalizes the results about dihedral groups from [4].

Theorem 2.1. *Let H and K be finite groups with bases such that there is a homomorphism $\phi : H \to \mathrm{Aut}(K)$. If $G = K \rtimes_\phi H$, then G has a basis.*

Proof. Let $(h_1, \dots, h_n)$ be a basis for H and $(k_1, \dots, k_m)$ be a basis for K. Then $h_1^{a_1} \cdots h_n^{a_n} = h_1^{c_1} \cdots h_n^{c_n}$ for some $a_i, c_i \in \{0, \dots, |h_i| - 1\}$ if and only if $(a_1, \dots, a_n) = (c_1, \dots, c_n)$ since $(h_1, \dots, h_n)$ is a basis for H, $(h_1^{a_1} \cdots h_n^{a_n})(h_1^{c_1} \cdots h_n^{c_n})^{-1} \in K$ if and only if $h_1^{a_1} \cdots h_n^{a_n} = h_1^{c_1} \cdots h_n^{c_n}$ since $H \cap K = \{(e, e)\}$, and $|G : K| = |H| = |h_1| \cdots |h_n|$. Therefore, the $(n+m)$-tuple $(k_1, \dots, k_m, h_1, \dots, h_n)$ is a basis for G by Lemma 1.2. □

The following corollary holds because A and B have bases as a consequence of the Fundamental Theorem of Abelian Groups. In particular, every dihedral group has a basis.

Corollary 2.2. *Every finite group isomorphic to $A \rtimes_\phi B$ for abelian groups A and B and some map $\phi : B \to \mathrm{Aut}(A)$ has a basis.*

2.2. **Special cases of solvable groups.** Proposition 1.1 demonstrates that not every solvable group has a basis. Still, we can demonstrate some special cases when solvable groups have bases. Recall that a subgroup H of a finite group G is a *Hall π-subgroup* for a set of primes π if $|H|$ and $|G : H|$ are coprime and every prime divisor of $|H|$ is in π, and a result of Philip Hall [5, Theorem 8.9] states that every solvable G has a Hall π-subgroup for any set of primes π.

Theorem 2.3. *Let G be a finite solvable group. If every Sylow subgroup of G has a basis, then G has a basis.*

Proof. Let $\pi(G)$ denote the primes that divide $|G|$. We will proceed by induction. If $|\pi(G)| = 1$, then G is equal to its sole Sylow subgroup, and thus has a basis by assumption.

Now suppose $|\pi(G)| > 1$. Let $p \in \pi(G)$ and $\pi'(G) = \pi(G) - \{p\}$. Because G is solvable, G has a proper Hall $\pi'(G)$-subgroup H. By induction, H has a basis $(h_1, \dots, h_m)$. Let P be a Sylow p-subgroup of G; then P has a basis $(b_1, \dots, b_n)$ by assumption. It is easy to check that H and $(b_1, \dots, b_n)$ fulfill the hypotheses of Lemma 1.2, so G has a basis. □

Recall that an integer m is *cube-free* if p^3 fails to divide m for every prime p.

Corollary 2.4. *Let m be an odd integer that is cube-free. If G has order m or $2m$, then G has a basis.*

Proof. Let p be a prime number that divides $|G|$, and let P be a Sylow p-subgroup of G. Then P must have order at most p^2 because $|G|$ is cube-free, which implies that P is abelian and thus has a basis. If $|G| = m$, then G is solvable by the Odd Order Theorem [1]. If $|G| = 2m$, then G is solvable by the Odd Order Theorem, [5, Corollary 6.12], and [5, Corollary 8.4]. Theorem 2.3 then implies G has a basis. □

3. Open Questions

We close with several open questions about group bases.

(1) We have seen that Q_8 does not have a basis. When exactly does a nilpotent group have a basis?
(2) Similarly, when does a solvable group have a basis? When does a non-solvable group have a basis?

References

1. W. Feit and J.G. Thompson, *Solvability of groups of odd order*, Pacific J. Math. **13** (1963), 775–1029.
2. M. I. González Vasco and R. Steinwandt, *Obstacles in two public key cryptosystems based on group factorizations*, Tatra Mt. Math. Publ. **25** (2002), 23–37.
3. P. Hall, *A contribution to the theory of groups of prime-power order*, Proceedings of the London Mathematical Society **2** (1934), no. 1, 29–95.
4. C. Hopkins, *Concerning uniqueness-bases of finite groups with applications to p-groups of class 2*, Transactions of the American Mathematical Society **41** (1937), no. 2, 287–313.
5. I.M. Isaacs, *Algebra: a graduate course*, Graduate Studies in Mathematics, vol. 100, American Mathematical Society, Providence, RI, 2009, Reprint of the 1994 original.
6. W. Lempken, T. van Trung, S. S. Magliveras, and W. Wei, *A public key cryptosystem based on non-abelian finite groups*, J. Cryptology **22** (2009), no. 1, 62–74.
7. S. S. Magliveras, *A cryptosystem from logarithmic signatures of finite groups*, Proceedings of the 29th Midwest Symposium on Circuits and Systems (1986), 972–975.
8. S. S. Magliveras, D. Stinson, and T. van Trung, *New approaches to designing public key cryptosystems using one-way functions and trapdoors in finite groups*, J. Cryptology **15** (2002), no. 4, 285–297.
9. P. Svaba and T. van Trung, *Public key cryptosystem* MST_3*: cryptanalysis and realization*, J. Math. Cryptol. **4** (2010), no. 3, 271–315.